



Supplementary Figure S1. Overview of *SILIA*-based phosphoproteomics.

The red and blue arrows show the workflow of two steps of our work. First, two sets of same seeds grown under identical conditions but labeled by distinct isotope of nitrogen is used to estimate incorporation rate of heavy nitrogen and construct standard curve (the red boxes), which will be used in the following calculation, following the red arrows. And the blue arrows show the *SILIA* approach to study the differentials between experimental set and control set quantitatively and the significantly changed phosphopeptides are determined (the blue box). A pair of both Forward and Reciprocal heavy nitrogen isotope labeling is always performed. For example, in the Forward experiment of a short-term ethylene-treated *rcn1-1*, the ethylene- and air-treated *rcn1-1* was labeled with ¹⁵N and ¹⁴N stable isotope-coded salt, respectively. Conversely, in the case of Reciprocal experiment, the two groups of plants were subjected to the nitrogen stable isotope labeling in a reverse order *i.e.* ¹⁴N for ethylene-treated and ¹⁵N for air-treated *rcn1-1*. This pair of labeling protocol is to eliminate the stable isotope labeling-resulting difference in ion intensity between ¹⁴N/¹⁵N-coded phosphopeptides (Kline *et al.*, 2010).




Supplementary Figure S2. False discovery rate (FDR) calculated using target-decoy searching.

Varying E -value (expectation value) is employed as the composite score for quality of peptide identification offered by the Mascot search engine. Red and blue curves show FDR estimated for identification of either ^{14}N - or ^{15}N -coded peptides and that for peptides discovered in pairs, *i.e.* both the ^{14}N - and ^{15}N -coded identical peptides varying with increase of E -value. The range around FDR = 1% is shown enlarged in the inset figure.



Supplementary Figure S3. Normalization of log₂ ratio of phosphopeptides.

Here m_r is the median value of all log₂ ratio, which is used to estimated actual mix ratio q using $m_r = \log_2(1/q)$. The media value m_r are subtracted from the log₂ ratio of each pair of phosphopeptide. All phosphopeptides detected in each biological replicate (either Forward or Reciprocal one) were quantified. The median values of log₂ ratios of all phosphopeptides from four mixtures of 1-min ethylene- and air-treated *rcn1-1* plants were found as 0.057, 0.183, -0.086 and 0.277 as well as that from four mixtures of wild type and *ctr1-1* mutants were found as 0.538, 0.287, 0.015 and 0.489. The ratio of each phosphopeptide quantified in each replicates were normalized based on the m_r value of the replicate.




Supplementary Figure S4. MS/MS spectrum of significantly altered phosphopeptides.

(A) and (B) MS/MS spectra of one-minute ethylene up- and down-regulated phosphopeptides, respectively.

(C) and (D) MS/MS spectrum of CTR1 up- and down-regulated phosphopeptides, respectively.

Both b-type ions and y-type ions are labeled to determine the peptide sequence and to localize the sites of phosphorylation. The singly and doubly H_3PO_4 neutral loss peptide ions are indicated as -98 and -196 in MS/MS spectrogram, respectively.




Supplementary Figure S5. Mass spectra of phosphopeptides enhanced by ethylene 1-min treatment in *rcn1-1*.

(A) TPNSDAEVIAL_pSPK from Indeterminate-Domain 5 (IDD5, AT2G02070);

(B) GLDIDTAGHHY_pTV from H⁺-ATPase 1 (HA1, AT2G18960);


(C) SL_pSDSEKAPYVAK from High Mobility Group B2 (HMGB2, AT1G20693), and/or High Mobility Group B3 (HMGB3, AT1G20696).

The phosphorylated residues are marked by *p*. Ethylene- and air-treatment are annotated by “E” and “A”, respectively.



Supplementary Figure S6. Mass spectra of phosphopeptides repressed by ethylene 1-min treatment in *rcn1-1*.

(A) SL_pSFSTSGPR from the leucine-rich repeat transmembrane protein kinase AT1G07650;
 (B) ATGAFILTA_pSHNPGGPTEDFGIK from Phosphoglucomutase 2 (PGM2, AT1G70730), and/or Phosphoglucomutase 3 (PGM3, AT1G23190);
 (C) GE_pSR_pSPPPYEKR from Arginine/Serine-rich Splicing Factor 41 (ATRSP41, AT5G52040).
 The phosphorylated residues are marked by *p*. Ethylene- and air-treatment are annotated by “E” and “A”, respectively.




Supplementary Figure S7. iTRAQ validations of significantly altered phosphorylation.

(A) and (B) MS spectra of iTRAQ-labeled phosphopeptides SL_pSDSEKAPYVAK (derived from HMGB2/B3, AT1G20693/AT1G20696) (A) and GE_pSR_pSPPPYEKR (derived from ATRSP41, AT5G52040) (B).

(C) and (D) MS/MS spectra of iTRAQ-labeled phosphopeptides SL_pSDSEKAPYVAK (C) and GE_pSR_pSPPPYEKR (D).

(E) and (F) Spectra of iTRAQ reporter ions detected for SL_pSDSEKAPYVAK (E) and GE_pSR_pSPPPYEKR (F) phosphorylated by kinase extracts from 0- (M/z 114), 1- (M/z 115), 5- (M/z 116) and 15-min (M/z 117) ethylene-treated plants.



Supplementary Figure S8. The MS spectrum of CTR1 up-regulated phosphopeptides.

(A) VQVA^pTVR from Plastid Transcriptionally Active 16 (PTAC16, AT3G46780);

(B) pSGTKFLPSSD from Calcium Sensing Receptor, (CaS, AT5G23060);


(C) GPSG^pSPWYGSDRVK from Light Harvesting Chlorophyll A/B Binding Protein 1.1 (LHCB1.1, AT1G29920), Light Harvesting Chlorophyll A/B Binding Protein 1.2 (LHCB1.2, AT1G29910), Light Harvesting Chlorophyll A/B Binding Protein 1.3 (LHCB1.3, AT1G29930), Light Harvesting Chlorophyll A/B Binding Protein 1.4 (LHCB1.4, AT2G34430), and/or Light Harvesting Chlorophyll A/B Binding Protein 1.5 (LHCB1.5, AT2G34420);

(D) KSA^pSFHTLDELEVR from the glutaredoxin family protein AT1G64500;

(E) GPSGSPWYG^pSDRVK from Light Harvesting Chlorophyll A/B Binding Protein 1.1 (LHCB1.1, AT1G29920), Light Harvesting Chlorophyll A/B Binding Protein 1.2 (LHCB1.2, AT1G29910), Light Harvesting Chlorophyll A/B Binding Protein 1.3 (LHCB1.3, AT1G29930), Light Harvesting Chlorophyll A/B Binding Protein 1.4 (LHCB1.4, AT2G34430), and/or Light Harvesting Chlorophyll A/B Binding Protein 1.5 (LHCB1.5, AT2G34420);

(F) oxMAPSIDAQLR from Phosphoenolpyruvate (PEP) carboxylase 2 (PPC2, AT2G42600), PEP carboxylase 3 (PPC3, AT3G14940), and/or PEP carboxylase-related protein AT3G42628.

The phosphorylated residues are marked by *p* and the oxidized Met is marked by *ox*. Wild type and *ctr1-1* mutant are annotated by “W” and “C”, respectively.



Supplementary Figure S9. The MS spectrum of CTR1 down-regulated phosphopeptides.

(A) SLG p SFR p SAANV from Plasma Membrane Intrinsic Protein 2A (PIP2A, AT3G53420), Plasma Membrane Intrinsic Protein 2B (PIP2B, AT2G37170), and/or Plasma Membrane Intrinsic Protein 2C (PIP2C, AT2G37180); (B) VG p SVQNWSK from the aluminium induced protein with YGL and LRDR motifs AT4G27450. The phosphorylated residues are marked by p . Wild type and *ctr1-1* mutant are annotated by “W” and “C”, respectively.